

A NOTE ON THE CONE OF MOBILE CURVES

MATEI TOMA

ABSTRACT. S. Boucksom, J.-P. Demailly, M. Păun and Th. Peternell proved that the cone of mobile curves $\overline{ME}(X)$ of a projective complex manifold X is dual to the cone generated by classes of effective divisors and conjectured an extension of this duality in the Kähler set-up. We show that their conjecture implies that $\overline{ME}(X)$ coincides with the cone of integer classes represented by closed positive smooth $(n-1, n-1)$ -forms. Without assuming the validity of the conjecture we prove that this equality of cones still holds at the level of degree functions.

Let X be a smooth complex projective variety of dimension n . A curve C on X is called *mobile* if it is the member of an algebraic family of (generically) irreducible curves covering X . Let $\overline{ME}(X)$ denote the closed convex cone generated by classes of mobile curves inside $N_1(X) := (H_{\mathbb{R}}^{n-1, n-1}(X) \cap H^{2n-2}(X, \mathbb{Z})/\text{Tors}) \otimes_{\mathbb{Z}} \mathbb{R}$. We shall call the elements of $\overline{ME}(X)$ *mobile classes*.

In [BDPP] it is shown that the following cones in $N_1(X)$ coincide:

- (1) the cone $\overline{ME}(X)$ of mobile curves,
- (2) the cone $\mathcal{M}_{NS} := \mathcal{M} \cap N_1$,
where $\mathcal{M} \subset H_{\mathbb{R}}^{n-1, n-1}(X)$ is the closure of the convex cone generated by cohomology classes of currents of the type $\nu_*(\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{n-1})$ for Kähler forms $\tilde{\omega}_1, \dots, \tilde{\omega}_{n-1}$ on a modification $\nu : \tilde{X} \rightarrow X$ de X ,
- (3) the dual cone $(\mathcal{E}_{NS})^{\vee}$ of the cone $\mathcal{E}_{NS}$ of pseudo-effective divisors on X .

It was known from [Dem92] that $\mathcal{E}_{NS} = \mathcal{E} \cap NS_{\mathbb{R}}$, where $\mathcal{E}$ is the cone of classes of positive closed currents of type $(1, 1)$ and $NS_{\mathbb{R}}(X) := (H_{\mathbb{R}}^{1,1}(X) \cap H^2(X, \mathbb{Z})/\text{Tors}) \otimes_{\mathbb{Z}} \mathbb{R}$. In [BDPP] Conj. 2.3 it is further conjectured that the cones $\mathcal{M}$ and $\mathcal{E}$ are dual.

In this note we compare $\overline{ME}(X)$ to the closed convex cone $P^{n-1, n-1}$ in $H_{\mathbb{R}}^{n-1, n-1}(X)$ generated by closed positive smooth $(n-1, n-1)$ -forms. From the above statements it is clear that $\overline{P^{n-1, n-1}} \cap N_1(X) \subset \overline{ME}(X)$. The converse inclusion will follow from our arguments if we admit the conjecture

Date: August 5, 2009.

AMS Classification (2000): 32J25; Secondary: 32J13, 14C30.

of Boucksom, Demailly, Păun, Peternell. If not we still get an equality at the level of degree functions as follows.

Any mobile class α gives rise to a degree function

$$\deg_\alpha : \text{Pic}(X) \rightarrow \mathbb{R}, \quad \deg_\alpha L := c_1(L)\alpha$$

and further to a notion of stability of torsion-free sheaves on X which generalizes the classical case $\alpha = H^{n-1}$ for a class H of an ample divisor, cf. [CaPe]. On the other side we consider *semi-Kähler* metrics on X , i.e. such that their associated Kähler forms ω satisfy $d\omega^{n-1} = 0$. Such a metric gives likewise a degree function:

$$\deg_\omega : \text{Pic}(X) \rightarrow \mathbb{R}, \quad \deg_\omega L := c_1(L)[\omega^{n-1}].$$

Then we can state:

Theorem. *For any mobile class α in the interior of the mobile cone $\overline{ME}(X)$ there exists a semi-Kähler metric on X with associated Kähler form ω such that $\deg_\alpha = \deg_\omega$ on $\text{Pic}(X)$.*

By [LT] Cor. 5.3.9 this has the following consequence on the moduli space of stable vector bundles:

Corollary. *If α is a class in the interior of $\overline{ME}(X)$, then the smooth part of the moduli space of stable vector bundles with respect to α admits a natural Kähler structure.*

The following notation will be used:

We denote by $\mathcal{D}^{p,q}$ the space of currents of bidegree (p, q) (and bidimension $(n-p, n-q)$) on X . Let

$$V_d = V_d(X) := \{T \in \mathcal{D}_\mathbb{R}^{1,1} \mid dT = 0\} / dd^c \mathcal{D}_\mathbb{R}^{0,0},$$

$$V_{dd^c} = V_{dd^c}(X) := \{T \in \mathcal{D}_\mathbb{R}^{1,1} \mid dd^c T = 0\} / \{\bar{\partial}S + \partial\bar{S} \mid S \in \mathcal{D}^{1,0}\},$$

$V_d^+, V_{dd^c}^+$ the cones generated by positive currents in V_d and V_{dd^c} respectively and

$V_{d,NS}^+, V_{dd^c,NS}^+$ their intersections with $NS_\mathbb{R}(X) := (H_\mathbb{R}^{1,1}(X) \cap H^2(X, \mathbb{Z})/\text{Tors}) \otimes_\mathbb{Z} \mathbb{R}$.

We first prove a

Lemma. *The natural map $j : V_d \rightarrow V_{dd^c}$ induces a bijection between positive cones*

$$V_{d,NS}^+ \rightarrow V_{dd^c,NS}^+.$$

Proof. It is known that in the case of compact Kähler manifolds the map j , which associates to a class of a bidegree $(1, 1)$ closed current $\{T\}_d \in V_d$ its class $\{T\}_{dd^c} \in V_{dd^c}$, is well defined and bijective.

The inclusion $j(V_d^+) \subset V_{dd^c}^+$ being obvious, we consider a dd^c -closed positive current T with $\{T\}_{dd^c} \in V_{dd^c, NS}$. Let $\eta := j^{-1}(\{T\}_{dd^c}) \in V_d$. We shall show that $\eta \in V_d^+$.

By the cited result of [BDPP] it suffices to check that for any modification $\nu : \tilde{X} \rightarrow X$ and Kähler forms $\tilde{\omega}_1, \dots, \tilde{\omega}_{n-1}$ on $\tilde{X}$ one has

$$\eta\nu_*([\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{n-1}]) \geq 0.$$

But

$$\eta\nu_*([\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{n-1}]) = \nu^*(\eta)[\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{n-1}] = \nu^*(\{T\}_{dd^c})[\tilde{\omega}_1 \wedge \dots \wedge \tilde{\omega}_{n-1}]$$

and Theorem 3 of [AlBa95] asserts that $\nu^*(\{T\}_{dd^c}) \in V_{dd^c}^+(\tilde{X})$, whence the desired inequality. $\square$

We can now give the proof of the Theorem.

Proof. Let α be an element in the interior of the cone of mobile curves. We denote by $\mathcal{D}_+^{1,1}$ the cone of positive currents inside the space $\mathcal{D}^{1,1}$ of bidegree $(1, 1)$ currents on X . We fix a Kähler form σ on X and set $\mathcal{D}_{+, \sigma}^{1,1} := \{T \in \mathcal{D}_+^{1,1} \mid \int_X T \wedge \sigma^{n-1} = 1\}$. This is a compact set for the weak topology on $\mathcal{D}^{1,1}$, [D] III.1.23. Let $\beta_1, \dots, \beta_k \in H_{\mathbb{R}}^{n-1, n-1}(X)$ be such that $V_{dd^c, NS} = \{t \in V_{dd^c} \mid t\beta_1 = 0, \dots, t\beta_k = 0\}$ and set $W := \{T \in \mathcal{D}^{1,1} \mid dd^c T = 0, \{T\}_{dd^c} \alpha = 0, \{T\}_{dd^c} \beta_1 = 0, \dots, \{T\}_{dd^c} \beta_k = 0\}$. Remark that W and $\mathcal{D}_{+, \sigma}^{1,1}$ are disjoint. Indeed, if $T \in \mathcal{D}_{+, \sigma}^{1,1}$ were dd^c -closed and $\{T\} \in V_{dd^c, NS}^+$, then by the Lemma there would exist a d -closed positive current $S \in \mathcal{D}_+^{1,1}$ such that $\{T\}_{dd^c} = j(\{S\}_d)$ and in particular

$$\{T\}_{dd^c} \alpha = \{S\}_d \alpha > 0.$$

The Hahn-Banach theorem then implies the existence of a functional on $\mathcal{D}^{1,1}$, which vanishes on W and is positive on $\mathcal{D}_{+, \sigma}^{1,1}$. This functional is thus given by a real $(n-1, n-1)$ -form u on X . The form u is strictly positive on X since the functional is positive on $\mathcal{D}_{+, \sigma}^{1,1}$. The vanishing on W implies that u is also d -closed. As functionals on $NS_{\mathbb{R}}(X)$, $[u]$ and α have the same kernel, hence they coincide up to some multiplicative constant.

It is enough now to take a positive $(n-1)$ -st root ω of u . For the convenience of the reader we show how this is done. Remark first that

$$(1), \quad (i \sum_{1 \leq i, j \leq n} a_{ij} dz_i \wedge d\bar{z}_j)^{n-1} = (n-1)! i^{(n-1)^2} \sum_{1 \leq i, j \leq n} (-1)^{i+j} c_{ji} \hat{d}z_i \wedge \hat{d}\bar{z}_j,$$

where we denoted by c_{ij} the cofactor of a_{ij} in the matrix $A = (a_{ij})_{1 \leq i, j \leq n}$ and $\hat{d}z_i := dz_1 \wedge \dots \wedge dz_{i-1} \wedge dz_{i+1} \wedge \dots \wedge dz_n$, $\hat{d}\bar{z}_j := d\bar{z}_1 \wedge \dots \wedge d\bar{z}_{j-1} \wedge d\bar{z}_{j+1} \wedge \dots \wedge d\bar{z}_n$. The relation ${}^tCA = \det(A)I_n$ for the matrix of cofactors $C = (c_{ij})_{1 \leq i, j \leq n}$ implies

$$A = {}^{n-1}\sqrt{\det(C)} {}^tC^{-1}$$

when A is positive definite. When the matrix C is positive definite, one obtains an unique positive definite solution A of equation (1). □

REFERENCES

- [AlBa95] ALESSANDRINI, L., BASSANELLI, G.: Modifications of compact balanced manifolds, *C. R. Acad. Sci. Paris Sr. I Math.* 320(1995), 1517–1522
- [BDPP] BOUCKSOM S., DEMAILLY J.-P., PĂUN M., PETERNELL TH. : The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension, *arXiv:math/0405285*
- [CaPe] CAMPANA F., PETERNELL TH. (WITH AN APPENDIX BY MATEI TOMA): Geometric stability of the cotangent bundle and the universal cover of a projective manifold, *arXiv:math/0405093*
- [Dem92] DEMAILLY J.-P: Regularization of closed positive currents and intersection theory. *J. Algebraic Geom.* 1 (1992), 361–409
- [D] DEMAILLY J.P.: Complex analytic and algebraic geometry, <http://www-fourier.ujf-grenoble.fr/~demailly/books.html>
- [LT] LÜBKE, M., TELEMAN, A. The Kobayashi-Hitchin correspondence, *World Scientific Publishing Co., Inc., River Edge, NJ, 1995*

INSTITUT DE MATHÉMATIQUES ELIE CARTAN, NANCY-UNIVERSITÉ,, B.P. 239, 54506
VANDOEUVRE-LÈS-NANCY CEDEX, FRANCE